



THE RESPONSE OF SOME CROP PLANTS TO PHYTOHORMONES AND SELECTED ELEMENTS

DISSERTATION

SUBMITTED IN PARTIAL FULFILMENT OF THE REQUIREMENTS

FOR THE AWARD OF THE DEGREE OF

Master of Philosophy

IN

BOTANY

SHAZIA ALVI

**DEPARTMENT OF BOTANY
ALIGARH MUSLIM UNIVERSITY
ALIGARH (INDIA)**

2001



28 JUL 2003



*This dissertation is humbly
dedicated to the memory of
my father*

DR.AQIL AHMAD
M.Phil.,Ph.D.
PD.Res.Train (Denmark)



PLANT PHYSIOLOGY SECTION
DEPARTMENT OF BOTANY
ALIGARH MUSLIM UNIVERSITY
ALIGARH-202002, U.P., INDIA
Phone: 91-571-408009
Fax: 91-571-702016
E-mail: aqil_ahmad@lycos.com

Dated: 24.9.2001

CERTIFICATE

This is to certify that **MS.SHAZIA ALVI** has worked under my supervision for the M.Phil. degree in Botany. She has fulfilled all conditions required to supplicate the M.Phil. degree. I therefore, approve that she may submit her dissertation entitled "*The response of some crop plants to phytohormones and selected elements*".

A handwritten signature in black ink, appearing to be "AQIL AHMAD", written over a horizontal line.

(DR.AQIL AHMAD)

Supervisor of Research

ACKNOWLEDGEMENT

In the name of Allah the most beneficent and the most merciful.

I avail the opportunity to express my profound sense of gratitude to my supervisor **Dr. Aqil Ahmad**, Reader, Department of Botany, who devoted much of his time going through the chapters, proof reading and finally looking after the printing of this dissertation. Without his help, this dissertation which had been long delayed, would not have seen the light of the day.

I am beholden to **Prof. Samiullah**, Chairman, Department of Botany, for his care and concern for the department, providing me the necessary facilities whenever I stood in need of from time to time.

I shall remain indebted to my other teachers, Drs. Arif Inam, Firoz Mohammad, Nafees A. Khan and M.M.A. Khan.

My thanks, in particular, are due to Dr. Shamsul Hayat who has contributed significantly in bringing this out speedily. My sincere thanks are also due to my colleagues, whose names are too many to be individually mentioned here, who have helped in the work in numerous ways.

I shall be failing in my duty, if I do not make a special mention of my father Late Mujtaba Ahmad Alvi, who always provided me with every facility regarding my studies and was anxious to see my work finished. Unfortunately, he passed away on 16th March 2001, yet I feel it was only due to his blessings that I could be able to finish my work.


(SHAZIA ALVI)

CONTENTS

S.NO.		Page No.
1.	Chapter -1 : Introduction	1 - 4
2.	Chapter -2 : Review of Literature	5 - 10
3.	Chapter -3 : Materials and Methods	11 - 16
4.	Chapter -4 : Results	17 - 18
5.	Chapter -5 : Discussion	19 - 21
6.	Bibliography	22 - 31
7.	Appendix	32 - 34

Chapter-1

Introduction

INTRODUCTION

The seed plays a unique role in the life history of the plant. The success with which the new individual is established is largely determined by the physiological and biochemical features of the seed. The seed which develops from the fertilized ovule is usually comprised of (1) the testa - the product of one or both integuments of the ovule; (2) the perisperm-derived from the nucellus; (3) the endosperm, produced as a result of the fusion between one male generative nucleus and the two polar nuclei to form the triploid endosperm nucleus and (4) the embryo, the result of fertilization of the oospore by a male nucleus.

Seeds characteristically contain large amount of food reserve which supports the growth and development of the seedling until it can establish itself as a photosynthesizing autotrophic plant. These reserves are laid down as discrete, intracellular bodies and include lipid, protein, carbohydrate, organic phosphate and various inorganic compounds. Some nutritionally undesirable reserves such as alkaloids, lectins, proteinase inhibitors, phytin and raffinose oligosaccharides are also present.

Germination is a series of events which take place when dry quiescent seeds imbibe water resulting in an increase in metabolic activity and the initiation of seedling from the embryo. The five stages involved in the seed germination and seedling growth are (1) imbibition and

hydration of protoplasm, (2) leakage of solutes, (3) activation and synthesis of enzymes (4) metabolism of storage products, their subsequent transport and synthesis of new materials and (5) cell elongation leading to the emergence of radicle.

Before germination process starts the seed takes up a large quantity of water. Many seeds placed under optimum favourable conditions for germination show a triphasic pattern of water uptake. Phase I occurs equally well in dead and living tissues and is, therefore, independent of the metabolic activity of the seed. Phase II is the period of active metabolism, in preparation for germination in non-dormant seeds of active metabolism in dormant seeds. Phase III is associated with germination and subsequent growth. During the initial stages of imbibition several substances like sugars, organic acids, phenols, inorganic phosphates and potassium leach out of the seed into the soil which might put the embryo under stress when the solubilization of stored food has just started. On hydration, during germination, the seed actively respire which supplies the energy (ATP); reducing power (NADH and NADPH) and carbon skeleton. Initially, there is a sharp increase in O_2 consumption, which can be partly attributed to the activation and hydration of mitochondrial enzymes involved in the krebs cycle and electron transport chain. The hydration of the seed is completed when oxygen uptake is stabilized and all pre-existing enzymes are activated. Shortly, following the absorption of water by the seed, enzyme activity begins to appear. Numerous enzymes such as lipases, proteinases, phosphatases, hydrolases, calmodulin, carboxypeptidases, and others (Bewly and Black 1985; Mayer and

Poljakoff-Mayber 1989; Coccuni and Negrini 1991; Washio and Ishikawa 1992; Bernier and Ballance 1993; Bernhardt *et al.* 1993) are either activated or synthesized a fresh. After the breakdown of the storage compounds into simpler chemical substances, they are transported from the endosperm cotyledons to the embryonic axis. These compounds are utilized in the production of new enzymes, structural materials, regulatory compounds, plant growth substances, and nucleic acids which carry out and regulate cell junctions and are consumed in the synthesis of new material. (Davies and Slack 1981; Mayer and Poljakoff-Mayber 1989).

The involvement of various plant hormones alone and in combination is based on the correlation of their endogenous concentration and the germination state. Gibberellins are a class of phytohormones which are most directly implicated in the germination process of the seeds. However, the involvement of the other major phytohormones (IAA, cytokinin, ABA, ethylene) cannot be over sighted. The role of auxins is not specifically assigned during the course of seed germination but it is actively involved (Bewley and Black, 1985).

The seeds are loaded with a sufficient quantity of cations and anions where calcium and potassium are not only involved as structural units of the membranes but also regulate its functioning. They also take active part in regulating cellular metabolism by controlling the activation and *de novo* synthesis of certain proteins (Bewley and Black, 1985). There is always a cumulative response of the cellular metabolism to the calcium and other factors (light and phytohormones). Therefore, the concentration of Ca^{2+} ,

at the level of the cytosol, may be altered for the signalling purposes by transiently opening the calcium channels in the plasma membrane or an intracellular membrane through the use of phytohormones.


The present study was aimed to elucidate the interactive effect of exogenously applied phytohormone (IAA) and the ions (Ca^{2+} and K^{+}) in regulating the seed metabolism at various intervals of seed germination to correlate their role.

Chapter-2

Review of Literature

REVIEW OF LITERATURE

S.No.		Page No.
2.1	Introduction	5-7
2.2	Metabolic changes associated with germination	7-9
2.3	Interaction effect of elements and phytohormones on the physiology of germinating seeds	9-10



REVIEW OF LITERATURE

2.1 Introduction

The hydration of the mature seed induces the synthesis/release of hydrolytic enzymes to initiate the solubilization of reserve food material. The simpler substances released are either respired to gain energy or absorbed in the synthesis of cellular constituents, at the level of the embryo, facilitating its development and growth. However, the whole process of germination is an accumulative expression of various factors, borned either within the seed or outside it. The level of hormonal reserves in the seed is one of the major regulators of the whole process of germination.

In a number of isolated cases, hormones are found to promote germination of the seeds. The exogenous application of GA improved the pre cent germination of the seeds of *Eleusine coracana*, Sesame, *Brassica juncea* (Mohanty and Sahoo, 1992), *Vigna mungo* (Baghel and Yadava, 1991; Sharma and Saran, 1992), *Vigna radiata* (Patel and Saxena, 1994). The involvement of the GA is further strengthen by the observation of Thomas (1989), who could overcome the inhibitory effect of GA biosynthesis inhibitors by subsequent addition of solution of GA₄ and GA₇ or the GA-active compound (1-3 chlorophthalimide). It was therefore, suggested that the biosynthesis of gibberellin is essential for breaking the dormancy in celery seeds.

The cytokinin is present in sufficient quantity in the developing seeds but its values decrease as the maturity reaches (Krishnamoorthy, 1993) and again the various forms increase as the germination advances (Saha and Sircar, 1996). Therefore, the seed treatment of green gram and black gram with kinetin enhanced germination (Patel and Saxena, 1994).

The mechanism involved in the action of ethylene, in seed germination, is not quite clear, however, in many species, exogenous ethylene or ethephon stimulates seed germination. Even ACC (1-aminocyclopropane-1-carboxylic acid) could improve seed germination but less efficiently (Kepczynski and Kepczynska, 1997). Similarly, auxins (IAA, NAA and IBA) are also recognized to effect the germination of the seeds of *Dolichos biflorus* (Nanda and Nanda, 1995), *V. radiata* and *V. mungo* (Baghel and Yadava, 1991; Patel and Saxena, 1994) in *Vigna mungo* and *Arachis hypogaea* (Manikanand and Hakim, 1999).

The contents of macronutrients differ qualitatively as well as quantitatively between different seeds (Bewly and Black, 1985), therefore, soaking the seeds of *V. mungo* in the extract of cow dung or 50 ppm solution of phosphorus for 12 hours improved germination percentage, compared with those soaked in water (Joseph, 1991). However, under saline (NaCl, KCl, KNO₃, CaCl₂ or CaNO₃) conditions the process of germination was either delayed or slowed down in the seeds of *Phaseolus mungo* (Sharma and Saran, 1994), wheat (Kamboh, *et al.*, 2000). Contrary to this, the hypochlorides of sodium (NaClO) and calcium (CaClO₂) favoured the germination of the orchid seeds (Miyoshi and Mii, 1998).

The response of the seeds to various trace elements varied. Vijaya and Punnuswamy (1997) treated the seeds of *Vigna mungo* and cowpea with 100 mg/kg of ZnSO_4 , MnSO_4 or Na_2Mo_4 singly or in various combinations. The treatment augmented the germination of the seeds of *V. mungo* but had no effect on cowpea. Similarly, the other cultivars gave a positive response to Mn but Hg (Nandi *et.al.*, 1995) and fluoride (Dhananjayanath *et.al.*, 1993) decreased/inhibited the germination.

2.2 Metabolic changes associated with germination

Germination is the process which culminates in the emergence of the radicle. Therefore, one has to concentrate on the phenomenon going on in the first few hours where the cell enlargement is the major event and is influenced by the auxins and other chemicals favouring overall germination of the seed. It has to be necessarily associated with the synthesis of the proteins, as wall components and critical enzymes required in the regulation of both metabolic and catabolic processes.

Nitrate reductase (NR) is a phytochrome mediated enzyme and its activity is facilitated by light. Out of the four forms of NR in *Sinapis alba* the appearance of NR_1 , NR_2 and NR_3 was strongly stimulated by red light whereas, NR_4 required the presence of continuous light (Schuster *et.al.*, 1989). Light also effects the rate of nitrate absorption by increasing the activity of NRA. It was confirmed by inhibiting the activity of NR by tungstate, that eventually decreased nitrate absorption (Rao and Rains, 1976). The increased activity of NR, in the presence of light/nitrate, and glutamine synthase (GS) is a coordinated appearance attributed to the

de novo synthesis of the enzymes proteins but GS level being slightly above that of NR. It appears to be the safety measures that the accumulation of ammonia may be prevented (Weber *et.al.*, 1990). Apart from light, substrate (nitrate) does have promotive effect on NR activity (Schuster *et.al.*, 1989). In addition to phytochrome and substrate, phytohormones profoundly effect the activity of NR. While nitrate maintained a steady level, irrespective of the treatment. The level of NRA was found to be much higher in the aerial leafy part than in the rooting zone of the pea cuttings, fed with an aqueous solution of IAA or IBA, during early rooting stage (Ahmad, 1988). Moreover, both the uptake and activity of NR did not depend on the endogenous auxins of pea cuttings in the first few days of rooting (Ahmad, 1994). The other hormones (GA_3 and kinetin) induced a significant increase in the activity of NR and NiR (Premabatidevi, 1998).

α -amylase is a Ca-containing protein, synthesised on the rough endoplasmic reticulum of the cell. GA_3 and ABA among the hormones, are known to regulate α -amylase production in aleurone layer cells of *Hordeum vulgare* seeds. GA_3 increased and abscisic acid (ABA) decreased Ca-flux into the endoplasmic reticulum and the amount of Ca that accumulated in the ER of barley aleurone cells (Bush *et.al.*, 1993). Fluctuations in the endogenous level of GA_1 and ABA in the germinating seeds of *Hordeum vulgare* confirmed the hypothesis that GA and ABA regulate the induction of α -amylase. The level of GA_1 was highest on the second day while that of ABA was lowest on the first day of the seed germination (Kobayoshi *et.al.*, 1995). The embryo contributes in the

Hormones cause influx of Ca^{2+} into the cells through specific Ca^{2+} channels in the plasmamembrane raising the cytosolic level of the ion, triggering the cellular response (Lehninger *et.al.*, 1993; Ching and Choi, 1998). Calcium is implicated in ABA induced thermotolerance in *Zea mays* (Ming *et.al.*, 1998), improved cold tolerance in low temperature induced injuries in *Oryza sativa* (Ying *et.al.*, 1996). Similarly it mediates the activation of NAD^+ kinase by involving IAA (Lixia *et.al.*, 1999). It has also been suggested by Yael and Weiss (1999) that there is a relationship between intercellular level of Ca^{2+} , calmodulin, protein phosphorylation and dephosphorylation with GA iduced gene expression including chalcone synthase in *Petunia corolla*. The biological responses to the interaction effect of the calcium and the hormone may be through an increase in the level of calmodulin (CaM) protein which is quite evident in case of aleurone layer cells incubated in $\text{GA}_3 + \text{CaCl}_2$ but was reduced by ABA (Schurink *et.al.*, 1996). Moreover, indole acetic acid also stimulated protein secretion by activating Ca^{++} channels of plasmalemma vesicles (Mendvedev *et.al.*, 1999).

Chapter-3

Materials & Methods

MATERIALS AND METHODS

S.No.		Page No.
3.1	Proposed study	11
3.2	Seeds	11
3.3	Experiment	11
3.4	Chemical analysis	12
3.4.1	Estimation of nitrate reductase activity	12
3.4.1.1	Colour development	13
3.4.2	Estimation of nitrate	13
3.4.2.1	Preparation f powder	13
3.4.2.2	Extraction and colour developoment	14
3.4.3	Estimation of proteins	14
3.4.3.1	Extraction	15
3.4.3.2	Colour development	15
3.4.4	Estimation of calcium and potassium	15
3.4.4.1	Digestion of powder	16
3.4.4.1.1	Estimation of calcium	16
3.4.4.1.2	Estimation of potasisum	16
3.5	Statistical analysis	16

MATERIALS AND METHODS

3.1 Proposed study

To achieve the objectives framed in chapter one, following studies were conducted to explore the response of seeds (*Cicer arietinum* cv. BG-256) to Indole-3-acetic acid (IAA), Ca^{2+} or K^{+} during the early stages of imbibition, under controlled conditions.

3.2 Seeds

The seeds of *Cicer arietinum* L. cv. BG-256 were procured from the National Seed Corporation Ltd. I.A.R.I. New Delhi. The healthy seeds were surface sterilized with 0.01% mercuric chloride solution followed by repeated washing with double distilled water (DDW).

3.3 Experiment

An experiment was conducted to study the interactive effect of IAA with Ca^{2+} or K^{+} .

(A) The surface sterilized seeds were soaked in distilled water and aqueous solutions of IAA (10^{-8}M), calcium chloride (4 or 8 μM of Ca^{2+}), potassium chloride (6 or 12 μM of K^{+}) to be used as control.

(B) The surface sterilized seeds were first allowed to imbibe in 10^{-8}M solution of IAA in sterilized petri plates for 6 hours and subsequently transferred, after being soaked with blotting paper, to different petri plates containing solutions of 4 and 8 μM of Ca^{2+} or 6 and 12 μM of K^{+} for

another six hours. These treated seeds were thoroughly washed with DDW to remove the adhering solutions and then transferred to sterilized petriplates containing cotton, moistened in DDW.

(C) Surface sterilized seeds were first allowed to imbibe in 4 and 8 μM of Ca^{2+} or 6 and 12 μM of K^{+} solutions for 6 hours and then transferred to 10^{-8} M solution of IAA for another 6 hours. These treated seeds were washed with DDW and transferred to sterilized petriplates containing cotton, moistened in DDW.

Seeds were allowed to germinate in BOD incubator at $28 \pm 2^{\circ}\text{C}$ and sampled at 12, 24 and 36 hours of the soaking (which also include the duration of treatment).

The following characteristics were studied in the seed.

1. Nitrate reductase activity.
2. Nitrate content
3. Calcium and Potassium contents.
4. Total protein content.

3.4 Chemical analysis

3.4.1 *Estimation of nitrate reductase activity*

Nitrate reductase activity (NRA) in the seed was estimated according to the method of Jaworski (1971). Biological sample was cut into small pieces of which 200 mg was weighed and transferred into polythene vial (25 ml). 2.5 ml of phosphate buffer (appendix 1.1) 0.5 ml

of 0.2M potassium nitrate (appendix 1.2) solution and 2.5 ml of 5% isopropanol were added. At least two drops of chloromphenicol were also added to check bacterial growth in the test sample. The vial was incubated in the BOD incubator at $30\pm 2^{\circ}\text{C}$ for 2 hours, in the dark.

3.4.1.1 Colour development

In a test tube 0.4 ml of test extract and 0.3 ml each of sulphanilamide (appendix 1.3) and naphthyl ethylene diamine hydrochloride (NED-HCl, appendix 1.4) were added. A pink colour appeared. The sample was left for 20-30 minutes for maximum colour development. It was diluted upto 5 ml with DDW and each sample was read at 540 nm using "Spectronic-20" colorimeter. Calibration, for 100% transmittance, was done by using a blank consisting of 4.4 ml DDW and 0.3 ml each of sulphanilamide and NED-HCl.

Standard curve was plotted using graded concentrations of potassium nitrite. Optical density of the test extract was read on the calibrated curve and NRA was calculated in terms of $\text{nmoles NO}_2 \text{ g}^{-1}\text{h}^{-1} \text{FW}$.

3.4.2 Estimation of nitrate

The nitrate content was estimated following the method of Johnson and Ulrich (1950).

3.4.2.1 Preparation of powder

The samples were dried, over night, in an oven at 80°C . The grinding was done in an electric grinder and powder passed through a 72

mesh screen and stored in polythene vials for the estimation of nitrate. Each sample, before analysis, was again dried over night in an oven at 80°C on a clean sheet of paper. In the morning the samples were placed in a dessicator for 15 minutes for cooling under dry conditions.

3.4.2.2 Extraction and colour development

50 mg of the above powder was weighed and transferred to a dried centrifuge tube (25 ml) with the addition of 400 mg of calcium sulphate and 12.5 ml of DDW. The sample was centrifuged for 10 minutes, at 6000 rpm. the supernatant was transferred to a 50 ml conical flask containing 1 ml of 0.5% calcium carbonate suspension. The excess solution was evaporated on water-bath leaving the final volume to 5 ml only. 0.5 ml of H₂O₂ (30%) was added to the above solution and the flask was closed with a lid. After 5 minutes the lid was removed and the solution in the flask was heated further to dryness on a water-bath in order to remove the peroxide. The flask was cooled and 1.25 ml of phenol-di-sulphonic acid (appendix 2.1) was rapidly added with continuous stirring. 35 ml of distilled water was also added to this solution. Lastly, 3 ml of 50% ammonium hydroxide solution was pipetted into it, yellow colour developed. The transmittance was noted, after 15 minutes at 400 nm using a "Spectronic-20" colorimeter. A standard curve was plotted using the graded concentrations of standard potassium nitrate solution.

3.4.3 Estimation of proteins

Total protein level in the sample was determined following the method of Lowry *et.al.* (1951).

3.4.3.1 Extraction

50 mg of the oven dried powder was transferred to a mortar. Samples was grind with the addition of 1 ml of trichloroacetic acid (5%). It was transferred to a centrifuge tube with repeated washings and the final volume was made up to 5 ml with trichloroacetic acid. The material was centrifuged at 4000 rpm for 15 minutes and the supernatant was discarded. 5 ml of 1N sodium hydroxide (NaOH) was added to the residue. It was left for 30 minutes in a waterbath at 60°C. After cooling for 15 minutes, the mixture was centrifuged at 4,000 rpm for 15 minutes and the supernatant was collected in 25 ml volumetric flask. Volume was made up to the mark with the help of 1N NaOH and used for the estimation of total protein.

3.4.3.2 Colour development

One ml of NaOH extract was transferred to a test tube and 5 ml of reagent D (appendix 3.1) was added to it. The solution was mixed well and allowed to stand at room temperature for 15 minutes. 0.5 ml of folin phenol reagent (appendix 3.2) was added rapidly with immediate mixing. The blue colour developed. The per cent transmittance of this solution was read at 660 nm using 'spectronic-20' colorimeter. A blank was run with each sample. The total protein content was calculated by comparing the optical density of each sample with a calibration curve plotted by taking known graded dilutions of a egg albumin.

3.4.4 Estimation of calcium and potassium

The following methodology was followed for the estimation of calcium and potassium in the samples.

3.4.4.1 Digestion of powder

100 mg of dry powder of the sample was taken in a 50 ml kjeldhal flask. To this 2 ml of concentrated sulphuric acid was added and the mixture was heated on a digestion rack, for 2 hours. The contents turned black in this duration. After cooling for 15 minutes, 0.5 ml of chemically pure H_2O_2 (30%) was added drop by drop till a clear solution was obtained. The peroxide digested material was transferred to a 100 ml volumetric flask with at least three washings with distilled water and the volume made upto the mark.

3.4.4.1.1 Estimation of calcium

Flame photometer was used to read the calcium content in the digested samples with the help of calcium filter. Per cent quantity of the element was calculated by comparing the readings on a standard curve plotted by using known dilution of calcium carbonate.

3.4.4.1.2 Estimation of potassium

Potassium content in the digested aliquot was directly estimated flame photometrically following the procedure explained by Singh (1988) by using potassium filter. A blank was run side by side. The reading was compared with a calibration curve plotted by using known dilutions of a standard potassium chloride solution.

3.5 Statistical analysis

The experimental data was analysed statistically following the procedures explained by Gomez and Gomez (1984).critical difference (C.D.) at the 5 % level of probability was calculated.

Chapter-4

Experimental Results

RESULTS

S.No.		Page No.
4.1	Nitrate reductase activity	17
4.2	Nitrate content	17
4.3	Protein content	18
4.4	Calcium content	18
4.5	Potassium content	18

RESULTS

4.1 Nitrate reductase activity

There is a consistent increase in the activity of nitrate reductase (NR) upto 36 hours of the imbibition (Table 1). The level of the enzyme increased significantly by the treatment. A maximum increase was recorded in the seeds soaked for 6 hours in 10^{-8} M of IAA only. The values were 50%, 95% and 77% more than the control at 12, 24 and 36 hours of the germination. Treating the seeds with either of the ions (Ca^{2+} , K^{+}) alone or in association with the auxin did not prove as good as the auxin alone. However, soaking the seeds first in IAA followed in the ionic solution for 6 hours each, proved better than otherwise. Within the two ions tested, Ca^{2+} proved better than K^{+} in association with IAA. The best combination being IAA+ Ca^{++} ($8\text{ }\mu\text{m}$) where the values increased by 42%, 60% and 46% at the three samplings (12, 24 and 36 hours).

4.2 Nitrate content

The content of nitrate decreased linearly as the germination progressed (Table 2). The seeds soaked in water (control) had maximum level of the ion at all the stages of smapling. In all the cases where the seeds were treated with auxin alone or in any combination with Ca^{2+} or K^{+} , the nitrate concentration was significantly less than the control. However, the IAA (10^{-8}) alone was most effective in reducing (metabolizing) the nitrate level in the seeds.

Table 1. The interaction effect of IAA (10^{-8}M) with Ca^{2+} (4 and 8 μM) or K^{+} (6 and 12 μM) on the nitrate reductase activity ($\text{nmol NO}_3\text{h}^{-1}\text{g}^{-1}\text{FW}$) in the seeds of *Cicer arietinum* L. cv. BG-256, at 12, 24 and 36 hours of the sampling.

Treatments	Sampling time (Hours)		
	12	24	36
Control	200.11	288.40	359.03
IAA 10^{-8}M	300.17	565.03	635.66
Ca4 μM	247.20	341.37	405.10
Ca8 μM	268.35	383.45	441.49
K6 μM	229.54	305.45	369.25
K12 μM	258.97	349.30	410.15
Ca4 + IAA	260.35	396.65	471.30
Ca8 + IAA	274.13	443.45	510.25
K6 + IAA	251.45	370.13	440.35
K12 + IAA	263.66	405.10	460.29
IAA + Ca4	279.35	419.30	479.36
IAA + Ca8	284.68	461.25	525.14
IAA + K6	263.45	386.10	451.39
IAA + K12	269.38	429.35	485.25
C.D. at 5%	20.26	18.65	31.25

Table 2. The interaction effect of IAA (10^{-8}M) with Ca^{2+} (4 and 8 μM) or K^{+} (6 and 12 μM) on the nitrate content (ppm) in the seeds of *Cicer arietinum* L. cv. BG-256, at 12, 24 and 36 hours of the sampling.

Treatments	Sampling time (Hours)		
	12	24	36
Control	2.49	2.20	1.66
IAA 10^{-8}M	2.11	1.72	1.09
Ca4 μM	2.40	2.14	1.56
Ca8 μM	2.31	2.05	1.43
K6 μM	2.45	1.98	1.36
K12 μM	2.42	2.08	1.46
Ca4 + IAA	2.29	1.96	1.28
Ca8 + IAA	2.25	1.89	1.19
K6 + IAA	2.30	1.92	1.23
K12 + IAA	2.24	1.83	1.16
IAA + Ca4	2.22	1.79	1.21
IAA + Ca8	2.20	1.88	1.13
IAA + K6	2.18	1.83	1.18
IAA + K12	2.24	1.93	1.28
C.D. at 5%	0.16	0.21	0.29

4.3 Protein content

Total protein content exhibited a progressive increase with the advancement of germination (Table 3). All the treatments had a significant impact on the level of the protein, as compared with the water soaked, control. Throughout the germination period, the maximum protein content was recorded in the seeds pre-soaked in IAA (10^{-8}M) alone for 12 hours or 6 hours in IAA subsequently followed for 6 hours in Ca^{2+} (4 or 8 μM). The respective increase, over the control, is 23%, 15.75% and 17%, respectively. It was closely followed by K^{+} (12 μM) used alone or in association with IAA.

4.4 Calcium content

The per cent calcium content increased as the germination advanced (Table 4). All the treatments significantly increased the level of the ion, over the control. There was a parallel relationship between the calcium content of the seed and its contents in the soaking medium, with or without the presence of the auxin. Pre-soaking in potassium was most effective if the seeds were already treated with auxin.

4.5 Potassium content

The seeds possessed more potassium as the germination progressed and exhibited a significant response to the treatment (Table 5). There was a proportionate relationship between the potassium content of the seed as that of its concentration in the bathing medium. The ionic content was maximum in the seeds pre-treated with IAA (10^{-8}M) followed by 12 μM of potassium for 6 hours each. The calcium alone or in association with auxin had a comparable effect on the level of the potassium.

Table 3. The interaction effect of IAA (10^{-8}M) with Ca^{2+} (4 and 8 μM) or K^{+} (6 and 12 μM) on the protein content (%) in the seeds of *Cicer arietinum* L. cv. BG-256, at 12, 24 and 36 hours of the sampling.

Treatments	Sampling time (Hours)		
	12	24	36
Control	18.53	20.24	21.43
IAA 10^{-8}M	22.79	25.69	27.65
Ca4 μM	21.58	24.48	26.23
Ca8 μM	21.19	24.36	26.20
K6 μM	20.68	23.80	25.10
K12 μM	22.44	25.60	29.30
Ca4 + IAA	21.45	23.85	24.93
Ca8 + IAA	21.68	24.15	25.90
K6 + IAA	21.25	23.90	25.35
K12 + IAA	22.13	25.03	27.02
IAA + Ca4	23.00	25.85	27.80
IAA + Ca8	23.09	26.25	29.89
IAA + K6	21.68	24.68	26.45
IAA + K12	22.68	25.32	27.25
C.D. at 5%	0.98	0.79	1.35

Table 4. The interaction effect of IAA (10^{-8}M) with Ca^{2+} (4 and 8 μM) or K^{+} (6 and 12 μM) on the calcium content (%) in the seeds of *Cicer arietinum* L. cv. BG-256, at 12, 24 and 36 hours of the sampling.

Treatments	Sampling time (Hours)		
	12	24	36
Control	0.125	0.132	0.137
IAA 10^{-8}M	0.139	0.143	0.147
Ca4 μM	0.156	0.159	0.163
Ca8 μM	0.169	0.174	0.179
K6 μM	0.143	0.149	0.154
K12 μM	0.145	0.151	0.156
Ca4 + IAA	0.168	0.174	0.178
Ca8 + IAA	0.170	0.188	0.191
K6 + IAA	0.148	0.151	0.156
K12 + IAA	0.147	0.153	0.162
IAA + Ca4	0.158	0.170	0.178
IAA + Ca8	0.180	0.181	0.196
IAA + K6	0.153	0.159	0.163
IAA + K12	0.152	0.157	0.166
C.D. at 5%	0.011	0.013	0.009

Table 5. The interaction effect of IAA (10^{-8}M) with Ca^{2+} (4 and 8 μM) or K^{+} (6 and 12 μM) on the potassium content (%) in the seeds of *Cicer arietinum* L. cv. BG-256, at 12, 24 and 36 hours of the sampling.

Treatments	Sampling time (Hours)		
	12	24	36
Control	0.532	0.546	0.553
IAA 10^{-8}M	0.565	0.573	0.580
Ca4 μM	0.568	0.577	0.583
Ca8 μM	0.564	0.573	0.580
K6 μM	0.603	0.613	0.624
K12 μM	0.629	0.636	0.641
Ca4 + IAA	0.570	0.583	0.596
Ca8 + IAA	0.569	0.579	0.586
K6 + IAA	0.613	0.626	0.633
K12 + IAA	0.639	0.647	0.656
IAA + Ca4	0.579	0.589	0.597
IAA + Ca8	0.583	0.593	0.604
IAA + K6	0.636	0.651	0.663
IAA + K12	0.660	0.673	0.677
C.D. at 5%	0.021	0.018	0.023

Chapter-5

Discussion

DISCUSSION

The process of seed germination starts with the imbibition of water and is accompanied with an increase in the metabolic activity. It is induced by the activation and the *de novo* synthesis of the hydrolytic enzymes (Bewley and Black, 1985). A larger share of these simpler substances, so produced, is carried to the embryonic axis to sustain its growth (Beevers, 1968). However, the rate and the direction of catabolic and anabolic reactions may, to some extent, be modified by selected chemicals including phytohormones (Davies, 1987) and the inorganic elements (Marschner, 1986).

A perusal of the observations (Table 1), like that of Tahir and Farooq (1989), revealed an increase in the activity of nitrate reductase as the germination process advanced. However, the contents of the nitrate (Table 2) showed a reverse trend, in confirmity with Egami *et al.* (1957). The NR is an inducible enzyme. Out of the factors which regulate the level of NR include the presence or absence of the substrate (Hewitt and Afridi, 1959; Afridi and Hewitt, 1964) and the phytohormones (Roth-Bejerane and Lips, 1970; Ahmad 1994, Ahmad and Hayat, 1999; Hayat *et al.*, 2001). In the present study, the seeds were simply soaked in distilled water/ a specific concentration of the auxin and/or the inorganic ions ($\text{Ca}^{2+}/\text{K}^{+}$). The germinating seeds therefore, totally depended, for its nitrate requirement, on the reserves. Under these uniform conditions, the IAA and/or the

cations ($\text{Ca}^{2+}/\text{K}^+$) elevated the level of the enzyme in the germinating seeds of *Cicer arietinum*. The auxin alone proved to be most effective in inducing the activity of the enzyme by possibly involving the genes because the synthesis of this enzyme involves the synthesis of both mRNA and protein (Jones and Prasad, 1992). The increase in the cytosolic level of calcium is known to trigger the cellular response (Stryer, 1992). Therefore, one of the observed expressions of the increase (Tables 4 & 5) in either of the ions (K^+ or Ca^{2+}) is the elevated level of NR. The other enzyme (α -amylase) is known to have a direct association with the level of the Ca^{2+} , during germination of rice seed (Ying *et al.*, 1996). The impact of the K^+ on the level of NR is possibly because of its involvement as an activator, co-factor in about 46 enzymes (Evans and Sorger, 1966). The treatment of the seeds with the either of the ions before or after being fed with auxin proved fruitful but the values of NR activity were less than those generated by the auxin alone. This variation may be due to the difference in the soaking pattern because in the former the duration of auxin treatment was just the half of the latter.

The nitrate content (Table 2) decreased linearly with the advancement of the germination as it was not supplemented from outside in the soaking medium. However, the rate of the loss (reduction) varied with the treatment but maximum quantity was found in the control at each stage of sampling. The nitrate content and the NRA showed inverse relationship with each other. It is obvious from these observations that a larger part of the nitrate was reduced under high NR activity. Therefore,

NR may be an appropriate marker enzyme for defining the importance of reduced nitrogen in the seed to takeup and reduced nitrate to nitrogen.

The major rate limiting stage in the reduction of nitrate to ammonical nitrogen for being incorporated in the production of amino acids is the very first step where nitrate is reduced to nitrite and is regulated by the enzyme NR (Hopkins, 1995). It is, therefore, understood that the increase in the activity of NR by the treatment (Table 1) will obviously enhance the availability of organic nitrogen and in turn that of amino acids. It possibly facilitated the synthesis of specific enzymes required for the synthesis of proteins (Hopkins, 1995) where its level had a concomitant increase (Table 3) with an increase in the level of NR. There seems to be a parallel relationship between NR and the level of the protein which is also supported by Singh and Singh (1985). The possible sites where the hormone and the ions could have acted is either at the level of the membrane and/or that of the genes.

Bibliography

BIBLIOGRAPHY

- Afridi, M.M.R.K. and Hewitt, E.J. 1964. The inducible formation and stability of nitrate reductase in higher plants. Effect of molybdenum in enzyme activity in cauliflower (*Brassica oleracea*). *J. Exp. Bot.* 15 : 251.
- Ahmad, A. 1988. Nitrate accumulation and nitrate reductase activity during rooting of pea cuttings treated with auxins. *Ind. J. Exp. Biol.* 26 : 470-472.
- Ahmad, A., 1994. Shoot apex as a source of auxin for nitrate uptake and actiity of nitrate reductase in pea cuttings. *Ind. J. Exp. Biol.* 32 : 65-67.
- Ahmad, A. and Hayat, S. 1999. Response of nitrate reductase to substituted IAA in pea seedlings. "*Plant Physiology for sustainable Agriculture*" (Srivastava, G.C., Singh, K. and Pal, M. Eds). pp. 252-259. Pointer Publisher, Jaipur, India.
- Appleford, N.E.J. and Lenton, J.R. 1997. Hormonal regulation of a-amylase gene expression in gemminating wheat (*Triticum aestivum*) grains. *Physiol. Plant.* 100 : 534-542.
- Arora, S. and Dua, S. 1994. Chemo-stress induced biochemical changes during germination of rapeseed. *Plant Physiol. Biochem.* 21: 44-45.
- Baghel, M.S. and Yadava, H.S. 1991. Response of black gram to dates of sowing and growth regulators. *Bhartiya-Krishi-Anusandhan Patrika.* 6: 99-104.

- Beevers, L., Schrader, L.E., Flesher, D. and Hageman, R.H. 1968. The role of light in the induction of nitrate reductase in radish cotyledons and maize seedlings. *Plant Physiol.* 40: 691-698.
- Bernhardt, D., Trutwig, A. and Barkhold, A. 1993. Synthesis of DNA and the development of amylase and phosphatase activities in cotyledons of germinating seeds of *Vaccaria pyramidata*. *J. Exp. Bot.* 44: 695-699.
- Bernier, A.M. and Ballance, G.M. 1993. Induction and secretion of alpha-amylase, (1-3), (1-4)-beta-glucanase, and (1-3)-beta-glucanase activities in gibberellic acid and CaCl_2 -treated half seeds and aleurones of wheat. *Cereal Chem.* 70: 127-132.
- Bewly, J.D. and Black M. 1985. Seed physiology of Development and germination. Plenum Publishing Corporation, New York.
- Bhatia, D.S., Ralhan, N., Malik, C.P., Ralhan, N., Grover, I.S. and Thukral, K. 1998. Role of mixtalol in assimilate transport in germinating *Cicer arietinum* seeds. *Environ. Develop.* 23 : 251-255.
- Bisambaev, A.K., Tairov, M.M. and Bersimbaev, R.I. 1992. Role of protein synthesis and Ca^{2+} in the effect of gibberellic acid on alpha amylase secretion from isolated aleurone layers of wheat seeds. *Biochemistry New York, Publ.* 57: 1273-1277.
- Bush, D.S., Biswas, A.K. and Jones, R.L. 1993. Hormonal regulation of Ca^{2+} transport in the endomembrane system of the barley aleurone. *Planta* 189: 507-515.

- Ching Hee-Don and Young-June Choi. 1998. Effect of temperature and foliar application of NNA and GA₃ on ⁴⁵Ca translocation and accumulation in the grafted melon (*Cucumis melo* var. *makuwa*) plants. *J. Korean Soc. Hort. Sci.* 39: 491-496.
- Cocucci, M. and Negrini, N. 1991. Calcium calmodulin in germination of *Phacelia tanacetifolia* seeds : Effects of light, temperature, fusicoccin and calcium calmodulin antagonists. *Physiol. Plant.* 82: 143-149.
- Davies, H.V. and Slack, P.T. 1981. The control of food mobilization in seeds of dicotyledonous plants. *New Phytol.* 88: 41-51.
- Davies, P.J. 1987. Plant hormones and their role in plant growth and development. Martinus Nijhoff, Boston, U.S.A.
- Dhananjayanath, V., Karunagaran, D. and Rao, P.R. 1993. Effect of flouride on germination and starch mobilization of horse gram seeds. *Ind. J. Plant Physiol.* 36: 131-133.
- Egami, F., Ohmachi, K., Iida, K. and Taniguchi, S. 1957. *Biochemia* 22: 122.
- Evans, H.J. and Sorger, G.J. 1966. *Annu. Rev. Plant Physiol.* 17: 47-76.
- Gomez, K.A. and Gomez, A.A. 1984. Statistical procedure for agricultural research. 2nd Eds. John Wiley & Sons, New York.
- Hayat, S., Ahmad, A., Hussain, A. and Mobin, M. 2001. Growth of wheat seedlings raised from the grains treated with 28-homobrassinolide. *Acta Physiol. Plant.* 23 : 27-30.

- Hewitt, E.J. and Afridi, M.M.R.K. 1959. Adaptive synthesis of nitrate reductase in higher plants. *Nature* 183: 57-58.
- Hopkins, G.W. 1995. Plants and nitrogen. In *"Introduction to plant physiology"*, pp. 118. John Wiley and Sons, New York.
- Ishizawa, K. and Esashi, Y. 1988. Action mechanism of ethylene in the control of sugar translocation in relation of rice coleoptile growth I sucrose metabolism. *Plant Cell Physiol.* 29: 131-141.
- Jain, R., Chowdhry, L. and Chatterjee, C. 1997. Influence of calcium of growth, composition and yield of blackgram (*V. mungo* L.) *Ind. J. Physiol.* 2: 221-224.
- Johnson, C.M. and Ulrich, A., 1950. Determination of nitrate in plant material. *Anal. Chem.* 22: 1526-1531.
- Jones, A.M. and Prasad, P. 1992. Auxin binding proteins and their possible roles in auxin mediated plant cell growth. *Bioassays*, 14: 43-48.
- Joseph, K. 1991. Studies on the effect of soil moisture content and seed soaking on the germination of legumes. *Leg. Res.* 14: 153-154.
- Jaworski, E.G. 1971. Nitrate reductase assay in intact plant tissues. *Biochem. Biophysics Res. Commun.* 43: 1274-1279.
- Kamboh, Ahmad, M., Oki, Y. and Adachi, T. 2000. Effect of pre-sowing seed treatments on germination and early seedlings growth of wheat varieties under saline conditions. *Soil Sci. Plant Nutr.* 46: 249-255.

- Kepczynski, J. and Kepczynka, E. 1997. Ethylene in seed dormancy and germination. *Physiol. Plant.* 101: 720-726.
- Kobayashi, M., Saitama, Gomi, M., Agematsu, J., Asami, T., Yoshida, S. and Sakurai, A. 1995. Fluctuations of endogenous gibberellin and abscisic acid level in germinating seeds of barley (*Hordeum vulgare*). *Biosci. Biotech. Biochem.* 59 : 1969-1970.
- Krishnamoorthy, H.N. 1993. Physiology of plant growth and development. Atma Ram and Sons, Delhi.
- Lehninger, A.L., Nelson, D.L., Cox, M.M. 1993. Principles of Biochemistry. CBS Publishers and Distributors, New Delhi.
- Lixia, Wang Ning-Ning, Wang Shu-Fang and Zhu Liang-Ji 1999. The effect of IAA on NAD kinase in wheat coleoptile is mediated by Ca^{2+} /Calmodulin complex. *Acta Phytophysiol. Sinica* 25: 357-361.
- Lowry, O.H., Rosenbrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193: 265-75.
- Manikanand, R. and Hakim, S.A. 1999. Effect of IAA and parthenium crude extract on blackgram and groundnut. *Adv. Plant Sci.* 12: 345-348.
- Marschner, H. 1986. Mineral nutrition of higher plants. Academic Press, London.

- Mayer, A.M. and Poljakoff-Mayber, A. 1989. The germination of seeds. Fourth Edition, Pergamon Press, London.
- Mendvedev, S.S., Batov, A.Y., Moshkov, A.V. and Markova, I.V. 1999. The role of ion channels in transduction of the auxin signal. *Fiziologiya Rastenii* (Moscow) 46: 711-717.
- Ming, G., Jun Li-Y and Zhong Chen, S. 1998. Absciscic acid induced thermotolerance in maize seedlings in mediated by calcium and associated with antioxidant systems. *J. Plant Physiol.* 153: 488-496.
- Misra, N., Dwivedi, U.N. 1995. Carbohydrate metabolism during seed germination and seedling growth in greengram under saline stress. *Plant Physiol. Biochem.* 33: 33-38.
- Miyoshi, K. and Mii, M. 1998. Stimulatory effects of sodium and calcium hypochlorite pre-chilling and cytokinin on the germination of *Cyperidium macranthos* seed in vitro. *Physiol. Plant.* 102: 481-486.
- Mohammad, K.A., Itoh, K., Iwabuchi, S., Hori, H. and Mitsu, T., 2000. Possible involvement of phosphoinositide Ca^{2+} signalling in the regulation of alpha amylase expression and germination of rice seed (*Oryza sativa*). *Plant Cell Physiol.* 41: 399-407.
- Mohanty, S.K. and Sahoo, N.C. 1992. Effect of soaking period, seed size and growth regulators on imbibition and gemrination of seeds of some field crops. *Orissa J. Agric. Res.* 5: 30-35.

- Nanda, I.P. and Nanda, H.P. 1995. Seed germination and seedling growth of hrose gram (*Dolichos biflorus*) treated with IAA. *Neo Botanica*. 3: 13-16.
- Nandi, S., Bera, A.K. and Nandi, S. 1995. Effect of mercury and manganese on seed germination and seedling growth in blackgram. *Seed Res*. 23: 125-128.
- Patel, I. and Saxena, O.P. 1994. Screening of PGRs for seed tgreatment in green gram and black gram. *Ind. J. Plant Physiol*. 37: 206-208.
- Premabatidevi, R.K. 1998. Effect of IAA, GA₃ and kinetin on nitrate reductase and nitrile reductase in the leaves of a tree legume (*Parkia javanica* Merr.) *Ind. J. Plant Physiol*. 3: 97-101.
- Qui, Z.S., Ling, D.V. and Jie Y.S. 1998. Effect of exogenous hormones on endogenous ABA level with its relation to assimilate accumulation of soyabean seeds. *Acta Bot. Sinica* 40: 642-646.
- Rao, K.P. and Rains, D.W. 1976. Nitrate absorption by Barley. Influence of nitrate reductase activity. *Amer. J. Plant Physiol*. 57: 59-62.
- Reddy, A.S.N., Friedmann, M. and Poovaiah, B.W. 1988. Auxin-induced changes in protein synthesis in the abscission zone of bean explants. *Plant Cell Physiol*. 29: 179-183.
- Roth-Bejerano, Lips S.H. 1970. Hormonal regulation of nitrate reductase activity. *New Phytol*. 69: 165-167.

- Saha, S. and Sircar, P.K. 1996. Cytokinin profile during early phase of germination of gram seeds after application of benzylaminopurine. *Biol. Plant.* 38: 293-296.
- Sangavanam, V. 1992. Mancozeb induced biochemical changes in the seedlings of Bengal gram. *Adv. Plant Sci.* 5:268-173.
- Schurink, R.C., Chang, P.V. and Jones, R.L. 1996. Modulation of calmodulin mRNA and protein levels in barley aleurone. *Plant Physiol.* 111: 371-380.
- Schuster, C., Schmidt, S. and Mohr H. 1989. Effect of nitrate ammonium light and a plastidic factor on the appearance of multiple forms of nitrate reductase in mustard (*Sinapis alba* L.) Cotyledons. *Planta* 177: 74-83.
- Sharma, A.K. and Saran, B. 1992. Effect of presowing soaking in NRA and GA₃ on gemrination and seedling growth in black gram. *New Agriculturist* 3: 21-24.
- Sharma, A.K. and Saran, B. 1994. Effect of salinity on germination and seedling growth in blackgram (*Phaseolus mungo* L.). *Neo-Botanica* 2: 57-59.
- Singh, L. 1988. Practical Agriculture Chemistry and Soil Science, published by Bishen Singh and Mahendra Pal Singh. Printed by Gajendra Singh Gahlot. Shiv Printers Dehradun.
- Singh, V.P. and Singh, M. 1985. Nitrate reductase activity and its relationship with grain protein and grain yield of wheat. *Ind. J. Plant Physiol.* . 28 : 235-242.

- Stryer, L. 1992. Biochemistry W.H. Freeman and Company New York.
- Tahir, I. and Farooq, S. 1989. Nitrate reductase activity in seedlings, leaves and grain of four buckwheats (*Fagopyrum* spp.) grown in Kashmir, India. *Acta Physiol. Plant* 11: 67-72.
- Thomas, T.H. 1989. Gibberellin involvement in dormancy break and germination of seeds of Celery *Apium graveolens* L. *Plant Growth Regul.* 8: 255-262.
- Torrent, M., Geli, M.I. and Ludevid, M.D. 1989. Storage protein hydrolysis and protein-body breakdown in germinated *Zea mays* L. seeds. *Planta* 180: 90-95.
- Tuan-Hua, David Ho, Abroms, J. and Varner J.E. 1982. Effect of ethylene on the release of alpha amylase through cell walls of Barley aleurone layers. *Plant Physiol.* 69: 1128-1131.
- Vijaya, A. and Ponnuswamy, A.S. 1997. Effect of seed fortification with micronutrients on seed quality in blackgram and cowpea. *Madras Agric. J.* 84: 223-224.
- Washio, K. and Ishikawa, K. 1992. Structure and expression during the germination of rice seeds of the gene for a carboxypeptidase. *Plant Mol. Biol.* 19: 631-640.
- Weber, M., Schmidt, S., Schuster, C. and Mohr, H. 1990. Factors involved in the coordinate appearance of nitrate reductase and glutamine synthase in the mustard (*Sinapis alba* L.) seedlings. *Planta* 180: 15-20.

- Yael, L.D. and Weiss, D. 1999. Ca^{2+} calmodulin and protein dephosphorylation are required for GA-induced gene expression in *Petunia corollas*. *Physiol. Plant.* **105**: 116-121.
- Ying, L., Sangen, W., Liang, L. and Wang, S.G. 1996. Effects of Ca^{2+} on seed vigour and cold resistance in rice (*Oryza sativa* L.) *J. Southwest Agric. Univ.* **18**: 491-495.
- Yu, W.Z., Zhang Fu-Suo, Wang, H.C. and Zou Chun Qin 1998. Effects of boron deficiency and low temperature on some physiological responses of cucumber seedlings. *Acta Phytophysiol. Sinica* **24**: 59-64.

Appendix

APPENDIX

PREPARATION OF REAGENTS

The various reagents used for biochemical determinations were prepared according to the following methods.

1.0 Reagents for the estimation of nitrate reductase activity

1.1 *Phosphate buffer (pH 7.5)*

- (a) 13.6 g potassium dihydrogen orthophosphate (KH_2PO_4) was dissolved in sufficient double distilled water and the final volume was adjusted upto 1 litre.
- (b) 17.42 g dipotassium monohydrogen orthophosphate (K_2HPO_4) was dissolved in sufficient double distilled water and the final volume was made upto 1 litre.
- (c) 160 ml of solution (a) and 840 ml of solution (b) was mixed in order to get pH 7.5.

1.2 *Potassium nitrate (0.2M)*

2.02 g potassium nitrate was dissolved in enough double distilled water and the final volume was made upto 100 ml.

1.3 *Sulphanilamide solution (1%)*

1 g sulphanilamide powder was dissolved in 100 ml 3N HCl.

1.4 NED-HCl solution (0.02%)

20 mg NED-HCl N-1-(naphthyl)-ethylene diamine dihydrochloric acid was dissolved in 100 ml double distilled water.

2.0 Reagent for the estimation of nitrate

2.1 Phenoldisulphonic acid reagent

25 g pure white phenol (AR) was dissolved in 150 ml of pure concentrated sulphuric acid to which 75 ml fuming sulphuric acid was added (13% SO_3). The solution was heated for about 2 hours at 100°C in water-bath and cooled. This solution was kept in a dark bottle in refrigerator.

3.0 Reagent for the estimation of protein

3.1 Reagent D

50 ml of 2% sodium carbonate was mixed with 1 ml of 0.5% copper sulphate and 1% sodium tartarate (which is prepared in the ratio of 1:1).

3.2 Folin's phenol reagent

100 g sodium tungstate and 25 g sodium molybdate was dissolved in 700 ml distilled water in which 50 ml of 85% phosphoric acid and 100 ml concentrated hydrochloric acid was mixed. The flask was connected with a reflux condenser and

boiled gently on a heating mantle for 10 hours. At the end of the boiling period, 150 g lithium sulphate, 50 ml double distilled water and 3-4 drops of liquid bromine was added to this flask. The reflux condenser was removed and the solution in the flask was boiled for 15 minutes in order to remove excess bromine, cooled and diluted to 1 litre.

The strength of this acid solution (1N) was tested by treating it with 1N sodium hydroxide using phenolphthalein as an indicator.

